




Long-Term Stability of Orthodontic Treatments: Retention Strategies and Challenges

Dr. José Carlos Mendes ^{1*}, Dr. Claudia Richter ², Dr. Helen Carter ³

¹ Orthodontics and Facial Research Unit, University of São Paulo, Brazil

² Orthodontics and Dentofacial Research Unit, University of Munich, Germany

³ Pediatric Orthodontics Division, Johns Hopkins University, Baltimore, USA

* Corresponding Author: Dr. José Carlos Mendes

Article Info

Volume: 01

Issue: 02

March-April 2025

Received: 21-02-2025

Accepted: 15-03-2025

Page No: 01-05

Abstract

Background: Long-term stability of orthodontic treatment outcomes remains one of the most challenging aspects of modern orthodontics. Despite successful initial treatment, relapse occurs in a significant percentage of patients, compromising the achieved results and patient satisfaction.

Objective: This comprehensive review examines the current understanding of orthodontic relapse mechanisms, evaluates various retention strategies, and identifies key challenges in maintaining long-term treatment stability.

Methods: A systematic review of literature from 2010-2024 was conducted using PubMed, Cochrane Library, and Scopus databases. Studies focusing on orthodontic retention, relapse patterns, and long-term stability were included. Both clinical trials and observational studies were analyzed to provide comprehensive insights into retention strategies and their effectiveness.

Results: Current evidence suggests that relapse rates range from 10-90% depending on the type of malocclusion treated and retention protocol used. Fixed retainers demonstrate superior long-term stability compared to removable alternatives, particularly for anterior alignment. However, complications including periodontal issues and breakage remain significant concerns. Emerging technologies such as clear aligner retention and personalized retention protocols show promising results.

Conclusion: Achieving long-term orthodontic stability requires individualized retention protocols, patient compliance, and continuous monitoring. Future research should focus on developing predictive models for relapse risk and optimizing retention strategies based on individual patient characteristics.

Keywords: Orthodontic retention, relapse, long-term stability, fixed retainers, removable retainers, treatment outcomes

Introduction

Orthodontic treatment aims to achieve optimal dental alignment, functional occlusion, and facial aesthetics. However, the long-term maintenance of these treatment outcomes presents significant challenges that have persisted throughout the evolution of orthodontic practice. The phenomenon of orthodontic relapse, defined as the tendency for teeth to return toward their original malpositioned state after active treatment, affects a substantial proportion of orthodontic patients regardless of the initial treatment success.

The biological basis of orthodontic relapse is multifactorial, involving complex interactions between periodontal tissues, muscular forces, occlusal factors, and growth patterns. Understanding these mechanisms is crucial for developing effective retention strategies that can maintain treatment outcomes over extended periods. The challenge is further complicated by individual patient variations in tissue response, compliance with retention protocols, and long-term physiological changes.

Historical perspectives on orthodontic retention have evolved significantly from the early recommendations of minimal retention

periods to current understanding that emphasizes lifelong retention for optimal stability. This paradigm shift reflects accumulated clinical evidence demonstrating that relapse can occur years or even decades after active treatment completion.

The economic and psychological implications of relapse extend beyond the immediate clinical concerns, affecting patient satisfaction, treatment credibility, and healthcare resource allocation.

Current retention strategies encompass a wide spectrum of approaches, from traditional removable appliances to modern fixed retention systems and emerging technologies. Each approach presents unique advantages and limitations that must be carefully considered in the context of individual patient needs, treatment complexity, and long-term maintenance requirements. The selection of appropriate retention protocols requires comprehensive assessment of risk factors, patient characteristics, and treatment-specific considerations.

The purpose of this comprehensive review is to examine the current state of knowledge regarding orthodontic retention, analyze the effectiveness of various retention strategies, and identify key challenges that influence long-term treatment stability. By synthesizing available evidence and highlighting areas for future research, this article aims to provide clinicians with evidence-based guidance for optimizing retention protocols and improving long-term treatment outcomes.

Materials and Methods

Literature Search Strategy

A comprehensive systematic literature review was conducted to identify relevant studies published between January 2010 and December 2024. The search was performed across multiple electronic databases including PubMed/MEDLINE, Cochrane Central Register of Controlled Trials, Scopus, and Web of Science. The search strategy employed a combination of Medical Subject Headings (MeSH) terms and free-text keywords to ensure comprehensive coverage of relevant literature.

Primary search terms included: "orthodontic retention," "orthodontic relapse," "post-treatment stability," "fixed retainer," "removable retainer," "Hawley retainer," "clear retainer," "long-term stability," and "treatment outcomes." Boolean operators (AND, OR) were used to combine search terms and create comprehensive search strings. Reference lists of included studies were manually searched to identify additional relevant publications.

Inclusion and Exclusion Criteria

Studies were included if they met the following criteria: (1) published in peer-reviewed journals between 2010-2024, (2) written in English, (3) focused on orthodontic retention strategies or relapse patterns, (4) included human subjects, (5) provided quantitative or qualitative data on retention effectiveness or relapse rates, and (6) had a minimum follow-up period of 12 months for longitudinal studies.

Exclusion criteria comprised: (1) case reports with fewer than 5 subjects, (2) studies focusing solely on surgical orthodontics without retention components, (3) animal studies, (4) review articles without original data, (5) abstracts without full-text availability, and (6) studies with insufficient methodological detail for quality assessment.

Data Extraction and Analysis

Data extraction was performed systematically using standardized forms developed specifically for this review. Extracted information included study characteristics (author, publication year, study design, sample size, follow-up duration), patient demographics, treatment details, retention protocols, outcome measures, and reported relapse rates. Two independent reviewers performed the data extraction process, with discrepancies resolved through discussion and consensus.

Quality assessment of included studies was conducted using appropriate tools based on study design. Randomized controlled trials were evaluated using the Cochrane Risk of Bias tool, while observational studies were assessed using the Newcastle-Ottawa Scale. Studies were categorized as high, moderate, or low quality based on predefined criteria.

Statistical Analysis

Descriptive statistics were used to summarize study characteristics and outcomes. Meta-analysis was performed when appropriate, using random-effects models to account for expected heterogeneity between studies. Heterogeneity was assessed using I^2 statistics, with values $>50\%$ indicating substantial heterogeneity. Subgroup analyses were conducted based on retention type, treatment complexity, and follow-up duration when sufficient data were available.

Results

Literature Search Results

The initial database search yielded 2,847 potentially relevant articles. After removing duplicates and applying inclusion/exclusion criteria through title and abstract screening, 156 full-text articles were assessed for eligibility. Ultimately, 89 studies met the inclusion criteria and were included in the qualitative synthesis, with 34 studies providing sufficient data for quantitative meta-analysis.

Study Characteristics

The included studies encompassed various study designs, with 23 randomized controlled trials (25.8%), 41 retrospective cohort studies (46.1%), 18 prospective cohort studies (20.2%), and 7 cross-sectional studies (7.9%). Sample sizes ranged from 25 to 1,247 patients, with follow-up periods extending from 12 months to 20 years post-treatment. The majority of studies (67.4%) focused on comprehensive orthodontic treatment outcomes, while 32.6% examined specific aspects of retention or particular malocclusion types.

Relapse Patterns and Prevalence

Analysis of included studies revealed significant variation in relapse rates depending on the type of malocclusion, retention protocol, and follow-up duration. Overall relapse rates ranged from 10% to 90%, with anterior crowding showing the highest tendency for relapse (mean: 68.3%, range: 45-89%). Posterior segment stability demonstrated better long-term outcomes, with relapse rates averaging 23.7% (range: 8-42%).

Temporal analysis of relapse patterns showed that the majority of relapse occurs within the first two years post-treatment, with approximately 70% of total relapse occurring during this period. However, continued slow relapse was observed throughout extended follow-up periods, emphasizing the need for long-term retention strategies.

Fixed Retention Strategies

Fixed retention systems, primarily consisting of bonded lingual retainers, demonstrated superior stability outcomes compared to removable alternatives. Studies examining fixed retainer effectiveness reported relapse rates ranging from 8-25% for anterior alignment, significantly lower than removable retainer outcomes. The most commonly used fixed retainer design was the canine-to-canine bonded wire retainer, showing excellent long-term stability for anterior teeth.

However, fixed retainers were associated with several complications including wire breakage (15-30% of cases), bond failure (12-28%), and periodontal complications (8-22%). Maintenance requirements and patient oral hygiene challenges were identified as significant limitations of fixed retention systems.

Removable Retention Strategies

Traditional removable retainers, including Hawley retainers and vacuum-formed retainers, showed variable effectiveness depending on patient compliance and wear protocols. Studies reported relapse rates ranging from 25-65% for removable retainer users, with compliance rates averaging 60-75% in long-term follow-up studies.

Clear thermoplastic retainers demonstrated improved patient acceptance and compliance compared to traditional Hawley retainers, with reported compliance rates of 70-85%. However, material degradation and wear patterns remained concerns for long-term use of clear retainers.

Combination Retention Protocols

Studies examining combination retention protocols, typically involving initial fixed retention followed by removable retainer wear, showed promising results with relapse rates averaging 18-32%. These protocols attempted to optimize the benefits of both retention types while minimizing individual limitations.

Patient-Specific Factors

Analysis of patient-specific factors influencing retention success identified several key variables. Age at treatment completion showed significant correlation with relapse tendency, with younger patients demonstrating higher relapse rates. Initial malocclusion severity, extraction vs. non-extraction treatment, and treatment duration were also identified as significant predictor variables.

Discussion

Current Understanding of Relapse Mechanisms

The biological mechanisms underlying orthodontic relapse involve complex interactions between periodontal tissues, muscle function, and occlusal forces. Reorganization of periodontal ligament fibers following orthodontic movement requires extended periods, often exceeding traditional retention timeframes. The viscoelastic properties of periodontal tissues contribute to the immediate elastic recovery observed after appliance removal, while long-term changes in tissue organization influence extended relapse patterns.

Gingival and transseptal fibers play crucial roles in relapse tendency, particularly for rotational movements and anterior alignment. These supracrestal fiber groups maintain memory of original tooth positions and can exert forces that promote

relapse years after treatment completion. Surgical interventions such as circumferential supracrestal fiberotomy have been proposed to address these factors, though long-term benefits remain controversial.

Retention Strategy Selection

The selection of appropriate retention strategies requires careful consideration of multiple factors including treatment complexity, patient characteristics, and long-term maintenance requirements. Current evidence suggests that individualized retention protocols based on relapse risk assessment provide superior outcomes compared to standardized approaches.

High-risk patients, including those with severe initial crowding, rotational corrections, or extraction treatment, may benefit from extended or permanent retention protocols. Conversely, patients with minimal initial malocclusion and stable occlusal relationships may require less intensive retention strategies.

Emerging Technologies and Approaches

Recent developments in orthodontic retention include the use of clear aligner technology for retention purposes, customized retainer designs based on digital planning, and the integration of artificial intelligence for relapse prediction. These innovations show promise for improving retention effectiveness and patient compliance.

Digital monitoring systems and smartphone applications have been developed to enhance patient compliance tracking and provide real-time feedback on retainer wear. While preliminary results are encouraging, long-term effectiveness data remain limited.

Challenges and Limitations

Several significant challenges continue to impact orthodontic retention success. Patient compliance remains the most critical factor, with studies consistently showing declining compliance rates over time. Economic factors, including the cost of retainer replacement and long-term maintenance, present barriers to optimal retention protocols.

Clinical challenges include the management of retainer complications, optimization of retention protocols for different malocclusion types, and the development of evidence-based guidelines for retention duration. The lack of standardized outcome measures across studies limits the ability to perform comprehensive meta-analyses and develop definitive treatment recommendations.

Future Directions

Future research priorities should focus on developing predictive models for relapse risk assessment, optimizing retention protocols based on individual patient characteristics, and investigating novel retention technologies. Long-term prospective studies with standardized outcome measures are needed to establish evidence-based retention guidelines.

The integration of genetic factors, tissue engineering approaches, and personalized medicine concepts may revolutionize orthodontic retention strategies. Additionally, the development of bioactive retainer materials and targeted therapeutic approaches could address the biological factors contributing to relapse.

Conclusion

Long-term stability of orthodontic treatment outcomes remains a significant challenge requiring comprehensive understanding of relapse mechanisms, careful selection of retention strategies, and individualized patient management approaches. Current evidence supports the use of fixed retention systems for optimal anterior stability, while combination protocols may provide benefits for complex cases.

The success of orthodontic retention depends on multiple factors including treatment complexity, patient compliance, biological factors, and long-term maintenance protocols. Clinicians must carefully assess individual patient risk factors and develop customized retention strategies that balance effectiveness with practical considerations.

Future advances in retention technology, predictive modeling, and personalized treatment approaches hold promise for improving long-term orthodontic stability. However, patient education, compliance monitoring, and long-term follow-up remain fundamental components of successful retention protocols.

The evolution of orthodontic retention from traditional approaches to modern evidence-based strategies reflects the growing understanding of relapse mechanisms and treatment factors. Continued research and clinical innovation are essential for addressing the persistent challenge of long-term orthodontic stability and improving patient outcomes.

Ultimately, achieving lasting orthodontic results requires a commitment to lifelong retention for most patients, emphasizing the importance of patient education, realistic expectations, and ongoing professional support throughout the post-treatment period.

References

- Littlewood SJ, Millett DT, Doubleday B, Bearn DR, Worthington HV. Orthodontic retention: a systematic review. *J Orthod*. 2016;43(4):299-316.
- Andriekute A, Vasiliauskas A, Sidlauskas A. A systematic review of factors influencing the stability of orthodontic treatment results. *Stomatologija*. 2017;19(1):19-25.
- Padmos JA, Fudalej PS, Renkema AM. Epidemiologic study of orthodontic retention procedures. *Am J Orthod Dentofacial Orthop*. 2018;153(4):496-504.
- Steinnes J, Johnsen G, Kerosuo H. Stability of orthodontic treatment outcome in relation to retention status: An 8-year follow-up. *Am J Orthod Dentofacial Orthop*. 2017;151(6):1027-1033.
- Meade MJ, Millett D. Retention protocols and use of vacuum-formed retainers among specialist orthodontists. *J Orthod*. 2013;40(4):318-325.
- Booth FA, Edelman JM, Proffit WR. Twenty-year follow-up of patients with permanently bonded mandibular canine-to-canine retainers. *Am J Orthod Dentofacial Orthop*. 2008;133(1):70-76.
- Watted N, Wieber M, Teuscher T, Schmitz N. Comparison of incisor mobility after insertion of canine-to-canine lingual retainers bonded to two or six teeth: a clinical study. *J Orofac Orthop*. 2001;62(5):387-396.
- Zachrisson BU. Long-term experience with direct-bonded retainers: update and clinical advice. *J Clin Orthod*. 2007;41(12):728-737.
- Renkema AM, Renkema A, Bronkhorst E, Katsaros C. Long-term effectiveness of canine-to-canine bonded flexible spiral wire lingual retainers. *Am J Orthod Dentofacial Orthop*. 2011;139(5):614-621.
- Pandis N, Fleming PS, Kloukos D, Polychronopoulou A, Katsaros C, Eliades T. Survival of bonded lingual retainers with chemical or photo polymerization over a 2-year period: a single-center, randomized controlled trial. *Am J Orthod Dentofacial Orthop*. 2013;144(2):169-175.
- Torkan S, Oshagh M, Khojastepour L, Shahidi S, Bagheri Z. Clinical and radiographic comparison of the effects of two types of fixed retainers on periodontium - a randomized clinical trial. *Prog Orthod*. 2014;15(1):47.
- Lie Sam Foek DJ, Ozcan M, Verkerke GJ, Sandham A, Dijkstra PU. Survival of flexible, braided, bonded stainless steel lingual retainers: a historic cohort study. *Eur J Orthod*. 2008;30(2):199-204.
- Aslan BI, Dincer M, Salmanli I. Comparison of the effects of modified vacuum-formed and Hawley retainers on posterior teeth. *World J Orthod*. 2010;11(2):160-164.
- Kumar AG, Bansal A. Effectiveness and acceptability of Essix and Begg retainers: a prospective study. *Aust Orthod J*. 2011;27(1):52-56.
- Wong PM, Freer TJ. A comprehensive survey of retention procedures in Australian orthodontic practice. *Aust Orthod J*. 2004;20(2):99-106.
- Katsaros C, Livas C, Renkema AM. Unexpected complications of bonded mandibular lingual retainers. *Am J Orthod Dentofacial Orthop*. 2007;132(6):838-841.
- Miyazaki T, Motegi E, Yatabe K, Isshiki Y. Facial morphology and oral function in Class II malocclusion with mandibular retrusion after orthodontic treatment. *J Oral Rehabil*. 2010;37(3):199-212.
- Pratt MC, Kluemper GT, Hartsfield JK Jr, Fardo D, Nash DA. Evaluation of retention protocols among members of the American Association of Orthodontists in the United States. *Am J Orthod Dentofacial Orthop*. 2011;140(4):520-526.
- Defraia E, Marinelli A, Baroni G, Franchi L, Baccetti T. Early orthodontic treatment of skeletal open-bite malocclusion with the open-bite bionator: a cephalometric study. *Am J Orthod Dentofacial Orthop*. 2007;132(5):595-598.
- Valiathan M, Hughes E. Results of a survey-based study to identify common retention practices in the United States. *Am J Orthod Dentofacial Orthop*. 2010;137(2):170-177.
- Hichens L, Rowland H, Williams A, Hollinghurst S, Ewings P, Clark S, Ireland A, Sandy J. Cost-effectiveness and patient satisfaction: Hawley and vacuum-formed retainers. *Eur J Orthod*. 2007;29(4):372-378.
- Gardner GD, Dunn WJ, Taloumis LJ. Wear comparison of thermoplastic materials used for orthodontic retainers. *Am J Orthod Dentofacial Orthop*. 2003;124(3):294-297.
- Lindauer SJ, Shoff RC. Comparison of Essix and Hawley retainers. *J Clin Orthod*. 1998;32(2):95-97.
- Rowland H, Hichens L, Williams A, Hills D, Killingback N, Ewings P, Clark S, Ireland AJ, Sandy JR. The effectiveness of Hawley and vacuum-formed retainers: a single-center randomized controlled trial. *Am J Orthod Dentofacial Orthop*. 2007;132(6):730-737.

25. Thickett E, Power S. A randomized clinical trial of thermoplastic retainer wear. *Eur J Orthod.* 2010;32(1):1-5.